Diabetologia

Clinical and Experimental Diabetes and Metabolism
Journal of the European Association for the Study of Diabetes (EASD)

Contents of Volume 46 • 2003

Editorials

Waldhäusl WK: Diabetologia and the Diabetologists 1
Waldhäusl WK: Diabetologia 1998–2003. Changing of the Guard 1591

Reviews

Chang C.K., Ulrich C.M.: Hyperinsulinaemia and hyperglycaemia: possible risk factors of colorectal cancer among diabetic patients 595
Fasshauer M., Paschke R.: Regulation of adipocytokines and insulin resistance 1594
Frühbeck G., Gómez-Ambrosi J.: Control of body weight: a physiologic and transgenic perspective 143
Gribble F.M., Reimann F.: Sulphonylurea action revisited: the post-cloning era 875
Kahn S.E.: The relative contributions of insulin resistance and beta-cell dysfunction to the pathophysiology of Type 2 diabetes 3
MacDonald P.E., Wheeler M.B.: Voltage-dependent K^+ channels in pancreatic beta cells: Role, regulation and potential as therapeutic targets 1046
Pyne N.J., Furman B.L.: Cyclic nucleotide phosphodiesterases in pancreatic islets 1179
Roep B.O.: The role of T-cells in the pathogenesis of Type 1 diabetes: From cause to cure 305
Rorsman P., Renström E.: Insulin granule dynamics in pancreatic beta cells 1029
Soltész G.: Diabetes in the young: a paediatric and epidemiological perspective 447
Taskinen M.-R.: Diabetic dyslipidaemia: from basic research to clinical practice 733
Yaney G.C., Corkey B.E.: Fatty acid metabolism and insulin secretion in pancreatic beta cells 1297
Yang Y., Santamaria P.: Dissecting autoimmune diabetes through genetic manipulation of non-obese diabetic mice 1447

Articles

Amoah A.G.B., Schuster D.P., Gaillard T., Osei K.: Insulin sensitivity and cardiovascular risk factors in hypertensive and normotensive native Ghanaians 949
Arroba A.I., Frago L.M., Pañeda C., Argente J., Chowen J.A.: The number of lactotrophs is reduced in the anterior pituitary of streptozotocin-induced diabetic rats (*Short Communication*) 634
Arteagoitia J.M., Larrañaga M.I., Rodriguez J.L., Fernandez I., Piniés J.A.: Incidence, prevalence and coronary heart disease risk level in known Type 2 diabetes: a sentinel practice network study in the Basque Country, Spain 899
Astill T.P., Ellis R.J., Arif S., Tree T.I.M., Peakman M.: Promiscuous binding of proinsulin peptides to Type 1 diabetes-permissive and -protective HLA class II molecules 496
Aye M., Sheedy W., Harrison R., Thompson J.S., Morice A.H., Masson E.A.: Pulmonary vasodilation in the rat by insulin in vitro could indicate potential hazard for inhaled insulin (Rapid Communication) 1199
Barberà A., Gudayol M., Eto K., Corominola H., Maechler P., Miró O., Cardellach F., Gomis R.: A high carbohydrate diet does not induce hyperglycaemia in a mitochondrial glycerol-3-phosphate dehydrogenase-deficient mouse 1394
Beltramo E., Buttiglieri S., Pomero F., Allione A., D'Alù F., Ponte E., Porta M.: A study of capillary pericyte viability on extracellular matrix produced by endothelial cells in high glucose 409
Boehm B.O., Lang G., Volpert O., Jehle P.M., Kurkhaus A., Rosinger S., Lang G.K., Bouck N.: Low content of the natural ocular anti-angiogenic agent pigment epithelium-derived factor (PEDF) in aqueous humor predicts progression of diabetic retinopathy 394
Bogdani M., Lefebvre V., Buelens N., Bock T., Pipeleers-Marichal M., In't Veld P., Pipeleers D.: Formation of insulin-positive cells in implants of human pancreatic duct cell preparations from young donors 830
Borg H., Arnqvist H.J., Björk E., Bolinder J., Eriksson J.W., Nyström L., Jeppsson J.-O., Sundkvist G.: Evaluation of the new ADA and WHO criteria for classification of diabetes mellitus in young adult people (15–34 yrs) in the Diabetes Incidence Study in Sweden (DISS) 173
Bos G., Dekker J.M., Nijpels G., Vegt F. de, Diamant M., Stehouwer C.D.A., Bouter L.M., Heine R.J.: A combination of high concentrations of serum triglyceride and non-high-density-lipoprotein-cholesterol is a risk factor for cardiovascular disease in subjects with abnormal glucose metabolism – The Hoorn Study 910
Bosco A.A., Lerario A.C., Santos R.F., Wajchenberg B.L.: Effect of thalidomide and rosiglitazone on the prevention of diabetic retinopathy in streptozotocin-induced diabetic rats 1669
Boulé N.G., Kenny G.P., Haddad E., Wells G.A., Sigal R.J.: Meta-analysis of the effect of structured exercise training on cardiorespiratory fitness in Type 2 diabetes mellitus 1071
Brennan L., Corless M., Hewage C., Malthouse J.P.G., McClenaghan N.H., Flatt P.R., Newsholme P.: ^{13}C NMR analysis reveals a link between L-glutamine metabolism, D-glucose metabolism and γ-glutamyl cycle activity in a clonal pancreatic beta-cell line 1512
Bui B.V., Armitage J.A., Tolcos M., Cooper M.E., Vingrys A.J.: ACE inhibition salvages the visual loss caused by diabetes 401
Burgdorf C., Richardt D., Kurz T., Adler S., Nötzold A., Kraatz E.G., Sievers H.H., Richardt G.: Norepinephrine release is reduced in cardiac tissue of Type 2 diabetic patients (*Short Communication*) 520
Burt D.J., Gruden G., Thomas S.M., Tutt P., Dell'Anna C., Viberti G.C., Gnudi L.: P38 mitogen-activated protein kinase mediates hexosamine-induced TGFβ1 mRNA expression in human mesangial cells 531

Cabrera-Rode E., Sarmiento L., Tiberti C., Molina G., Barrios J., Hernández D., Díaz-Horta O., Di Mario U.: Type 1 diabetes islet associated antibodies in subjects infected by echovirus 16 1348
Camp T.M., Tyagi S.C., Senior R.M., Hayden M.R., Tyagi S.C.: Gelatinase B(MMP-9) an apoptotic factor in diabetic transgenic mice 1438
Cardozo A.K., Proost P., Gysemans C., Chen M.-C., Mathieu C., Eizirik D.L.: IL-1β and IFN-γ induce the expression of diverse chemokines and IL-15 in human and rat pancreatic islet cells, and in islets from pre-diabetic NOD mice 255
Carlsson P.-O., Berne C., Östenson C.-G., Andersson A., Jansson L.: Hypoglycaemia induces decreased islet blood perfusion mediated by the central nervous system in normal and Type 2 diabetic GK rats 1124
Carvalheira J.B.C., Ribeiro E.B., Araújo E.P., Guimarães R.B., Telles M.M., Torsoni M., Gontijo J.A.R., Velloso L.A., Saad M.J.A.: Selective impairment of insulin signalling in the hypothalamus of obese Zucker rats 1629
Cattan P., Rottembourg D., Cottet S., Tardivel I., Dupraz P., Thorens B., Boitard C., Carel J.C.: Destruction of conditional insulinoma cell lines in NOD mice: A role for autoimmunity 504
Cederberg J., Picard J.J., Eriksson U.J.: Maternal diabetes in the rat impairs the formation of neural-crest derived cranial nerve ganglia in the offspring 1245
Chandie Shaw P.K., van Es L.A., Paul L.C., Rosendaal F.R., Souverijn J.H.M., Vandenbroucke J.P.: Renal disease in relatives of Indo-Asian Type 2 diabetic patients with end-stage diabetic nephropathy 618
Chang I., Cho N., Koh J.-Y., Lee M.-S.: Pyruvate inhibits zinc-mediated pancreatic islet cell death and diabetes 1220
Chang T.I., Horal M., Jain S.K., Wang F., Patel R., Loeken M.R.: Oxidant regulation of gene expression and neural tube development: Insights gained from diabetic pregnancy on molecular causes of neural tube defects 538
Choi J.B., Uchino H., Azuma K., Iwashita N., Tanaka Y., Mochizuki H., Migita M., Shimada T., Kawamori R., Watada H.: Little evidence of transdifferentiation of bone marrow-derived cells into pancreatic beta cells 1366
Cnop M., Havel P.J., Utzschneider K.M., Carr D.B., Sinha M.K., Boyko E.J., Retzlaff B.M., Knopp R.H., Brunzell J.D., Kahn S.E.: Relationship of adiponectin to body fat distribution, insulin sensitivity and plasma lipoproteins: evidence for independent roles of age and sex 459
Curtis T.M., Major E.H., Trimble E.R., Scholfield C.N.: Diabetes-induced activation of protein kinase C inhibits store-operated Ca^{2+} uptake in rat retinal microvascular smooth muscle 1252
Dabrowski M., Larsen T., Ashcroft F.M., Bondo Hansen J., Wahl P.: Potent and selective activation of the pancreatic beta-cell type K_{ATP} channel by two novel diazoxide analogues 1375
Davis B.J., Johnston C.I., Burrell L.M., Burns W.C., Kubota E., Cao Z., Cooper M.E., Allen T.J.: Renoprotective effects of vasopeptidase inhibition in an experimental model of diabetic nephropathy 961
De Breuck S., Lardon J., Rooman I., Bouwens L.: Netrin-1 expression in fetal and regenerating rat pancreas and its effect on the migration of human pancreatic duct and porcine islet precursor cells 926
De Souza C.T., Gasparetti A.L., Pereira-da-Silva M., Araújo E.P., Carvalheira J.B., Saad M.J.A., Boschero A.C., Carneiro E.M., Velloso L.A.: Peroxisome proliferator-activated receptor γ coactivator-1-dependent uncoupling protein-2 expression in pancreatic islets of rats: a novel pathway for neural control of insulin secretion 1522
de Vos P., Smedema I., van Goor H., Moes H., van Zanten J., Netters S., de Leij L.F.M., de Haan A., de Haan B.J.: Association between macrophage activation and function of micro-encapsulated rat islets 666
Delmeire D., Flamez D., Hinke S.A., Cali J.J., Pipeleers D., Schuit F.: Type VIII adenylyl cyclase in rat beta cells: coincidence signal detector/generator for glucose and GLP-1 1383
Deng S., Vatamaniuk M., Lian M.-M., Doliba N., Wang J., Bell E., Wolf B., Raper S., Matschinsky F.M., Markmann J.F.: Insulin gene transfer enhances the function of human islet grafts 386
Devendra D., Galloway T.S., Horton S.J., Evenden A., Keller U., Wilkin T.J.: The use of phage display to distinguish insulin autoantibody (IAA) from insulin antibody (IA) idiotypes 802
Diem P., Deplazes M., Fajfr R., Bearth A., Müller B., Christ E.R., Teuscher A.: Effects of alcohol consumption on mortality in patients with Type 2 diabetes mellitus 1581
Douek I.F., Gillespie K.M., Dix R.J., Bingley P.J., Gale E.A.M.: Three generations of autoimmune diabetes: an extended family study 1313
Eaton S.E.M., Harris N.D., Ibrahim S., Patel K.A., Selmi F., Radatz M., Ward J.D., Tesfaye S.: Increased sural nerve epineurial blood flow in human subjects with painful diabetic neuropathy 934
Edwards R., Burns J.A., McElduff P., Young R.J., New J.P.: Variations in process and outcomes of diabetes care by socio-economic status in Salford, UK 750
Eriksson J.G., Forsén T., Tuomilehto J., Osmond C., Barker D.J.P.: Early adiposity rebound in childhood and risk of Type 2 diabetes in adult life 190
Esberg L.B., Ren J.: Role of nitric oxide, tetrahydrobiopterin and peroxynitrite in glucose toxicity-associated contractile dysfunction in ventricular myocytes 1419
Escobar-Morreale H.F., Villuendas G., Botella-Carretero J.I., Sancho J., San Millán J.L.: Obesity, and not insulin resistance, is the major determinant of serum inflammatory cardiovascular risk markers in pre-menopausal women 625
Etoh T., Inoguchi T., Kakimoto M., Sonoda N., Kobayashi K., Kuroda J., Sumimoto H., Nawata H.: Increased expression of NAD(P)H oxidase subunits, NOX4 and p22phox, in the kidney of streptozotocin-induced diabetic rats and its reversibity by interventive insulin treatment 1428
Fanelli C.G., Pampanelli S., Porcellati F., Bartocci L., Scionti L., Rossetti P., Bolli G.B.: Rate of fall of blood glucose and physiological responses of counterregulatory hormones, clinical symptoms and cognitive function to hypoglycaemia in Type I diabetes mellitus in the postprandial state 53
Ferrannini E., Gastaldelli A., Miyazaki Y., Matsuda M., Pettiti M., Natali A., Mari A., DeFronzo R.A.: Predominant role of reduced beta-cell sensitivity to glucose over insulin resistance in impaired glucose tolerance 1211
Ferre T., Riu E., Franckhauser S., Agudo J., Bosch F.: Long-term overexpression of glucokinase in the liver of transgenic mice leads to insulin resistance 1662
Figarola J.L., Scott S., Loera S., Tessler C., Chu P., Weiss L., Hardy J., Rahbar S.: LR-90 a new advanced glycation endproduct inhibitor prevents progression of diabetic nephropathy in streptozotocin-diabetic rats 1140
Galeano M., Deodato B., Altavilla D., Cucinotta D., Arsic N., Marini H., Torre V., Giacca M., Squadrito F.: Adeno-associated viral vector-mediated human vascular endothelial growth factor gene transfer stimulates angiogenesis and wound healing in the genetically diabetic mouse 546
Gao G., Shao C., Zhang S.X., Dudley A., Fant J., Ma J.-X.: Kallikrein-binding protein inhibits retinal neovascularization and decreases vascular leakage (*Rapid Communication*) 689
Gardiner T.A., Anderson H.R., Degenhardt T., Thorpe S.R., Baynes J.W., Archer D.B., Stitt A.W.: Prevention of retinal capillary basement membrane thickening in diabetic dogs by a non-steroidal anti-inflammatory drug 1269
Gault V.A., O'Harte F.P.M., Harriott P.F., Mooney M.H., Green B.D., Flatt P.R.: Effects of the novel (Pro^3)GIP antagonist and exendin(9–39) amide on GIP- and GLP-1-induced cyclic AMP generation, insulin secretion and postprandial insulin

release in obese diabetic (*ob/ob*) mice: evidence that GIP is the major physiological incretin 222
Gerritsen J., TenVoorde B.J., Dekker J.M., Kingma R., Kostense P.J., Bouter L.M., Heethaar R.M.: Measures of cardiovascular autonomic nervous function: agreement, reproducibility, and reference values in middle age and elderly subjects 330
Gonzalez C., Cuvellier S., Hue-Beauvais C., Lévi-Strauss M.: Genetic control of non obese diabetic mice susceptibility to high-dose streptozotocin-induced diabetes 1291
Gross M.-L., El-Shakmak A., Szábó A., Koch A., Kuhlmann A., Münter K., Ritz E., Amann K.: ACE-Inhibitors but not endothelin receptor blockers prevent podocyte loss in early diabetic nephropathy 856
Gu D., Reynolds K., Duan X., Xin X., Chen J., Wu X., Mo J., Whelton P.K., He J., for the InterASIA Collaborative Group: Prevalence of diabetes and impaired fasting glucose in the Chinese adult population: International Collaborative Study of Cardiovascular Disease in Asia (InterASIA) 1190
Gual P., Grémeaux T., Gonzalez T., Le Marchand-Brustel Y., Tanti J.-F.: MAP kinases and mTOR mediate insulin-induced phosphorylation of Insulin Receptor Substrate-1 on serine residues 307, 612 and 632 1532
Guangda X., Yuhua W.: Apolipoprotein *e4* allele and endothelium-dependent arterial dilation in Type 2 diabetes mellitus without angiopathy 514
Gysemans C., Stoffels K., Giulietti A., Overbergh L., Waer M., Lannoo M., Feige U., Mathieu C.: Prevention of primary non-function of islet xenografts in autoimmune diabetic NOD mice by anti-inflammatory agents 1115
Hamilton J.S., Powell L.A., McMaster C., McMaster D., Trimble E.R.: Interaction of glucose and long chain fatty acids (C18) on antioxidant defences and free radical damage in porcine vascular smooth muscle cells in vitro 106
Hammarstedt A., Smith U.: Thiazolidinediones (PPARγ ligands) increase IRS-1, UCP-2 and C/EBPα expression, but not transdifferentiation, in L6 muscle cells 48
Hattori Y., Suzuki M., Hattori S., Kasai K.: Globular adiponectin upregulates nitric oxide production in vascular endothelial cells (*Rapid Communication*) 1543
Heijer T. den, Vermeer S.E., Dijk E.J. van, Prins N.D., Koudstaal P.J., Hofman A., Breteler M.M.B.: Type 2 diabetes and atrophy of medial temporal lobe structures on brain MRI 1604
Hermann R., Knip M., Veijola R., Simell O., Laine A.-P., Åkerblom H.K., Groop P.-H., Forsblom C., Pettersson-Fernholm K., Ilonen J., The FinnDiane Study Group: Temporal changes in the frequencies of HLA genotypes in patients with Type 1 diabetes – indication of an increased environmental pressure? 420
Hernandez R., Teruel T., Lorenzo M.: Rosiglitazone produces insulin sensitisation by increasing expression of the insulin receptor and its tyrosine kinase activity in brown adipocytes 1618
Hinck L., Thissen J.P., Pampfer S., De Hertogh R.: Effect of high concentrations of glucose on differentiation of rat trophoblast cells in vitro 276
Houard N., Rousseau G.G., Lemaigre F.P.: HNF-6-independent differentiation of mouse embryonic stem cells into insulin-producing cells 378
Houdali B., Wahl H.G., Kresi M., Nguyen V., Haap M., Machicao F., Ammon H.P.T., Renn W., Schleicher E.D., Häring H.-U.: Glucose oversupply increases Δ9-desaturase expression and its metabolites in rat skeletal muscle 203
Hu G., Qiao Q., Silventoinen K., Eriksson J.G., Jousilahti P., Lindström J., Valle T.T., Nissinen A., Tuomilehto J.: Occupational, commuting, and leisure-time physical activity in relation to risk for Type 2 diabetes in middle-aged Finnish men and women 322
Hu G., The DECODE Study Group: Gender difference in all-cause and cardiovascular mortality related to hyperglycaemia and newly-diagnosed diabetes 608
Idris I., Gray S., Donnelly R.: Rosiglitazone and pulmonary oedema: an acute dose-dependent effect on human endothelial cell permeability (*Rapid Communication*) 288
Inagaki Y., Yamagishi S., Okamoto T., Takeuchi M., Amano S.: Pigment epithelium-derived factor prevents advanced glycation end products-induced monocyte chemoattractant protein-1 production in microvascular endothelial cells by suppressing intracellular reactive oxygen species generation (*Short Communication*) 284
Iori E., Marescotti M.C., Vedovato M., Ceolotto G., Avogaro A., Tiengo A., Del Prato S., Trevisan R.: In situ protein Kinase C activity is increased in cultured fibroblasts from Type 1 diabetic patients with nephropathy 524
Izumi Y., Yamada K.A., Matsukawa M., Zorumski C.F.: Effects of insulin on long-term potentiation in hippocampal slices from diabetic rats 1007
Jellema A., Zeegers M.P.A., Feskens E.J.M., Dagnelie P.C., Mensink R.P.: *Gly972Arg* variant in the insulin receptor substrate-1 gene and association with Type 2 diabetes: a meta-analysis of 27 studies 990
Jonas J.C., Guiot Y., Rahier J., Henquin J.C.: Haeme-oxygenase 1 expression in rat pancreatic beta cells is stimulated by supraphysiological glucose concentrations and by cyclic AMP 1234
Kamiya J., Aoki Y.: Associations between hyperglycaemia and somatic transversion mutations in mitochondrial DNA of people with diabetes mellitus 1559
Kästenbauer T., Hörnlein B., Sokol G., Irsigler K.: Evaluation of granulocyte-colony stimulating factor (Filgrastim) in infected diabetic foot ulcers (Short Communication) 27
Kato Y., Miura Y., Yamamoto N., Ozaki N., Oiso Y.: Suppressive effects of a selective inducible nitric oxide synthase (iNOS) inhibitor on pancreatic beta-cell dysfunction 1228
Kawashima J., Tsuruzoe K., Motoshima H., Shirakami A., Sakai K., Hirashima Y., Toyonaga T., Araki E.: Insulin down-regulates resistin mRNA through the synthesis of protein(s) that could accelerate the degradation of resistin mRNA in 3T3-L1 adipocytes 231
Kefas B.A., Heimberg H., Vaulont S., Meisse D., Hue L., Pipeleers D., Van de Casteele M.: AICA-riboside induces apoptosis of pancreatic beta cells through stimulation of AMP-activated protein kinase (*Short Communication*) 250
Kim K.-A., Kang K., Chi Y.-I., Chang I., Lee M.-K., Kim K.-W., Shoelson S.E., Lee M.-S.: Identification and functional characterization of a novel mutation of hepatocyte nuclear factor-1α gene in a Korean family with MODY3 721
Kitamura T., Kimura K., Makondo K., Furuya D.T., Suzuki M., Yoshida T., Saito M.: Proinsulin C-peptide increases nitric oxide production by enhancing mitogen-activated protein-kinase-dependent transcription of endothelial nitric oxide synthase in aortic endothelial cells of Wistar rats 1698
Klimes I., Weston K., Kovacs P., Gasperikova D., Jezova D., Kvetnansky R., Thompson J.R., Sebokova E., Samani N.J.: Mapping of genetic loci predisposing to hypertriglyceridaemia in the hereditary hypertriglyceridaemic rat: analysis of genetic association with related traits of the insulin resistance syndrome 352
Koistinen H.A., Chibalin A.V., Zierath J.R.: Aberrant p38 mitogen-activated protein kinase signalling in skeletal muscle from Type 2 diabetic patients (*Rapid Communication*) 1324
Krebs M., Brehm A., Krssak M., Anderwald C., Bernroider E., Nowotny P., Roth E., Chandramouli V., Landau B.R., Waldhäusl W., Roden M.: Direct and indirect effects of amino acids on hepatic glucose metabolism in humans 917
Kriz J., Padjen A.L.: Intra-axonal recording from large sensory myelinated axons: Demonstration of impaired membrane conductances in early experimental diabetes 213
Krueger T., Wohlrab U., Klucken M., Schott M., Seissler J.: Autoantigen-specific protection of non-obese diabetic mice from

cyclophosphamide-accelerated diabetes by vaccination with dendritic cells 1357
Kukko M., Kimpimäki T., Kupila A., Korhonen S., Kulmala P., Savola K., Simell T., Keskinen P., Ilonen J., Simell O., Knip M.: Signs of beta-cell autoimmunity and HLA-defined diabetes susceptibility in the Finnish population: the sib cohort from the Type 1 Diabetes Prediction and Prevention Study 65
Kullin M., Li Z., Bondo Hansen J., Welsh N., Karlsson F.A., Sandler S.: Protection of rat pancreatic islets by potassium channel openers against alloxan, sodium nitroprusside and interleukin-1β mediated suppression – possible involvement of the mitochondrial membrane potential 80
Laing S.P., Swerdlow A.J., Slater S.D., Burden A.C., Morris A., Waugh N.R., Gatling W., Bingley P.J., Patterson C.C.: Mortality from heart disease in a cohort of 23,000 patients with insulin-treated diabetes 760
Lamont B.J., Andrikopoulos S., Funkat A., Favaloro J., Ye J.M., Kraegen E.W., Howlett K.F., Zajac J.D., Proietto J.: Peripheral insulin resistance develops in transgenic rats overexpressing phosphoenolpyruvate carboxykinase in the kidney 1338
Larsen M.O., Gotfredsen C.F., Wilken M., Carr R.D., Pørksen N., Rolin B.: Loss of beta-cell mass leads to a reduction of pulse mass with normal periodicity, regularity and entrainment of pulsatile insulin secretion in Göttingen minipigs 195
Lee D.-H., Ha M.-H., Kim J.-H., Christiani D.C., Gross M.D., Steffes M., Blomhoff R., Jacobs D.R. Jr: Gamma-glutamyltransferase and diabetes – a 4 year follow-up study 359
Lee Y.H., Tokraks S., Pratley R.E., Bogardus C., Permana P.A.: Identification of differentially expressed genes in skeletal muscle of non-diabetic insulin-resistant and insulin-sensitive Pima Indians by differential display PCR 1567
Lefrandt J.D., Bosma E., Oomen P.H.N., Hoeven J.H. van der, Roon A.M. van, Smit A.J., Hoogenberg K.: Sympathetic mediated vasomotion and skin capillary permeability in diabetic patients with peripheral neuropathy 40
Lindsay J.R., McKillop A.M., Mooney M.H., O'Harte F.P.M., Bell P.M., Flatt P.R.: Demonstration of increased concentrations of circulating glycated insulin in human Type 2 diabetes using a novel and specific radioimmunoassay 475
Looker H.C., Fagot-Campagna A., Gunter E.W., Pfeiffer C.M., Venkat Narayan K.M., Knowler W.C., Hanson R.L.: Homocysteine as a risk factor for nephropathy and retinopathy in Type 2 diabetes 766
Lopes C., Dina C., Durand E., Froguel P.: PAI-1 polymorphisms modulate phenotypes associated with the metabolic syndrome in obese and diabetic Caucasian population 1284
Luik P.T., Hoogenberg K., Kerstens M.N., Beusekamp B.J., Jong P.E. de, Dullaart R.P.F., Navis G.J.: The influence of the ACE (*I/D*) polymorphism on systemic and renal vascular responses to angiotensins in normotensive, normoalbuminuric Type 1 diabetes mellitus 1131
Lynn S., Borthwick G.M., Charnley R.M., Walker M., Turnbull D.M.: Heteroplasmic ratio of the A3243G mitochondrial DNA mutation in single pancreatic beta cells (*Short Communication*) 296
Malik R.A., Kallinikos P., Abbott C.A., van Schie C.H.M., Morgan P., Efron N., Boulton A.J.M.: Corneal confocal microscopy: a non-invasive surrogate of nerve fibre damage and repair in diabetic patients 683
Martin S., Kolb H., Beuth J., Leendert R. van, Schneider B., Scherbaum W.A.: Change in patients' body weight after 12 months of treatment with glimepiride or glibenclamide in Type 2 diabetes: a multicentre retrolective cohort study 1611
McCrimmon R.J., Jacob R.J., Fan X., McNay E.C., Sherwin R.S.: Effects of recurrent antecedent hypoglycaemia and chronic hyperglycaemia on brainstem extra-cellular glucose concentrations during acute hypoglycaemia in conscious diabetic BB rats (*Short Communication*) 1658
Meier J.J., Gallwitz B., Siepmann N., Holst J.J., Deacon C.F., Schmidt W.E., Nauck M.A.: Gastric inhibitory polypeptide (GIP) dose-dependently stimulates glucagon secretion in healthy human subjects at euglycaemia (*Short Communication*) 798
Mensink M., Blaak E.E., Vidal H., Bruin T.W.A. de, Glatz J.F.C., Saris W.H.M.: Lifestyle changes and lipid metabolism gene expression and protein content in skeletal muscle of subjects with impaired glucose tolerance 1082
Mitamura Y., Harada T., Harada C., Ohtsuka K., Kotake S., Ohno S., Tanaka K., Takeuchi S., Wada K.: NF-κB in epiretinal membranes after human diabetic retinopathy 699
Miyanaga F., Ogawa Y., Ebihara K., Hidaka S., Tanaka T., Hayashi S., Masuzaki H., Nakao K.: Leptin as an adjunct of insulin therapy in insulin-deficient diabetes 1329
Miyazawa-Hoshimoto S., Takahashi K., Bujo H., Hashimoto N., Saito Y.: Elevated serum vascular endothelial growth factor is associated with visceral fat accumulation in human obese subjects 1483
Mottagui-Tabar S., Rydén M., Löfgren P., Faulds G., Hoffstedt J., Brookes A.J., Andersson I., Arner P.: Evidence for an important role of perilipin in the regulation of human adipocyte lipolysis 789
Movassat J., Beattie G.M., Lopez A.D., Portha B., Hayek A.: Keratinocyte growth factor and beta-cell differentiation in human fetal pancreatic endocrine precursor cells 822
Murata G.H., Shah J.H., Adam K.D., Wendel C.S., Bokhari S.U., Solvas P.A., Hoffman R.M., Duckworth W.C.: Factors affecting diabetes knowledge in Type 2 diabetic veterans 1170
Nakabayashi M., Morishita R., Nakagami H., Kuba K., Matsumoto K., Nakamura T., Tano Y., Kaneda Y.: HGF/NK4 inhibited VEGF-induced angiogenesis in in vitro cultured endothelial cells and in vivo rabbit model 115
Nakamura M., Kawahara M., Morishige N., Chikama T., Nakata K., Nishida T.: Promotion of corneal epithelial wound healing in diabetic rats by the combination of a substance P-derived peptide ($FGLM\text{-}NH_2$) and insulin-like growth factor-1 (*Short Communication*) 839
Nannipieri M., Posadas R., Williams K., Politi E., Gonzales-Villalpando C., Stern M.P., Ferrannini E.: Association between polymorphisms of the Atrial Natriuretic Peptide gene and proteinuria: a population-based study (*Short Communication*) 429
Nayak R.C., Agardh C.D., Kwok M.G.K., Stjernquist H., Farthing-Nayak P.J., Agardh E.: Circulating anti-pericyte autoantibodies are present in Type 2 diabetic patients and are associated with non-proliferative retinopathy (*Short Communication*) 511
Niemi M., Backman J.T., Granfors M., Laitila J., Neuvonen M., Neuvonen P.J.: Gemfibrozil considerably increases the plasma concentrations of rosiglitazone (*Short Communication*) 1319
Niemi M., Backman J.T., Neuvonen M., Neuvonen P.J.: Effects of gemfibrozil, itraconazole, and their combination on the pharmacokinetics and pharmacodynamics of repaglinide: potentially hazardous interaction between gemfibrozil and repaglinide (*Rapid Communication*) 347
Nivoit P., Wiernsperger N., Moulin P., Lagarde M., Renaudin C.: Effect of glycated LDL on microvascular tone in mice: a comparative study with LDL modified in vitro or isolated from diabetic patients 1550
Ota T., Takamura T., Ando H., Nohara E., Yamashita H., Kobayashi K.: Preventive effect of cerivastatin on diabetic nephropathy through suppression of glomerular macrophage recruitment in a rat model 843
Pagetta A., Folda A., Brunati A.M., Finotti P.: Identification and purification from the plasma of Type 1 diabetic subjects of a proteolytically active Grp94. Evidence that Grp94 is entirely responsible for plasma proteolytic activity 996
Palm F., Cederberg J., Hansell P., Liss P., Carlsson P.-O.: Reactive oxygen species

cause diabetes-induced decrease in renal oxygen tension 1153
Panhuysen C.I.M., Cupples L.A., Wilson P.W.F., Herbert A.G., Myers R.H., Meigs J.B.: A genome scan for loci linked to quantitative insulin traits in persons without diabetes: the Framingham Offspring Study 579
Park S.-H., Park J.-W., Park S.-J., Kim K.-Y., Chung J.-W., Chun M.-H., Oh S.-J.: Apoptotic death of photoreceptors in the streptozotocin-induced diabetic rat retina 1260
Park Y., Freedman B.D., Lee E.J., Park S., Jameson J.L.: A dominant negative PPARγ mutant shows altered cofactor recruitment and inhibits adipogenesis in 3T3-L1 cells 365
Pedersen-Bjergaard U., Agerholm-Larsen B., Pramming S., Hougaard P., Thorsteinsson B.: Prediction of Severe Hypoglycaemia by Angiotensin-Converting Enzyme Activity and Genotype in Type 1 Diabetes 89
Pelikánová T., Šimková R., Tesař V., Jirsa M.: Effect of acute hyperglycaemia on selected plasma and urinary cytokine antagonists in Type 1 diabetes mellitus (*Short Communication*) 470
Perrin D., Soulage C., Pequignot J.M., Géloën A.: Resistance to obesity in Lou/C rats prevents ageing-associated metabolic alterations 1489
Poa N.R., Cooper G.J.S., Edgar P.F.: Amylin gene promoter mutations predispose to Type 2 diabetes in New Zealand Maori 574
Pomero F., Allione A., Beltramo E., Buttiglieri S., D'Alù F., Ponte E., Lacaria A., Porta M.: Effects of protein kinase C inhibition and activation on proliferation and apoptosis of bovine retinal pericytes 416
Popov D., Simionescu M., Shepherd P.R.: Saturated-fat diet induces moderate diabetes and severe glomerulosclerosis in hamsters 1408
Potashnik R., Bloch-Damti A., Bashan N., Rudich A.: IRS1 degradation and increased serine phosphorylation cannot predict the degree of metabolic insulin resistance induced by oxidative stress 639
Pouwer F., Beekman A.T.F., Nijpels G., Dekker J.M., Snoek F.J., Kostense P.J., Heine R.J., Deeg D.J.H.: Rates and risks for co-morbid depression in patients with Type 2 diabetes mellitus: results from a community-based study 892
Pöykkö S., Ukkola O., Kauma H., Savolainen M.J., Kesäniemi Y.A.: Ghrelin Arg51Gln mutation is a risk factor for Type 2 diabetes and hypertension in a random sample of middle-aged subjects (*Short Communication*) 455
Pruhova S., Ek J., Lebl J., Sumnik Z., Saudek F., Andel M., Pedersen O., Hansen T.: Genetic epidemiology of MODY in the Czech republic: new mutations in the MODY genes *HNF-4α*, *GCK* and *HNF-1α* (*Short Communication*) 291
Queen L.R., Ji Y., Goubareva I., Ferro A.: Nitric oxide generation mediated by β-adrenoceptors is impaired in platelets from patients with Type 2 diabetes mellitus 1474
Rasschaert J., Liu D., Kutlu B., Cardozo A.K., Kruhøffer M., Ørntoft T.F., Eizirik D.L.: Global profiling of double stranded RNA- and IFN-γ-induced genes in rat pancreatic beta cells 1641
Rathmann W., Haastert B., Icks A., Löwel H., Meisinger C., Holle R., Giani G.: High prevalence of undiagnosed diabetes mellitus in Southern Germany: Target populations for efficient screening. The KORA survey 2000 182
Reimann F., Huopio H., Dabrowski M., Proks P., Gribble F.M., Laakso M., Otonkoski T., Ashcroft F.M.: Characterisation of new K_{ATP}-channel mutations associated with congenital hyperinsulinism in the Finnish population 241
Renard C.B., Askari B., Suzuki L.A., Kramer F., Bornfeldt K.E.: Oleate, not ligands of the receptor for advanced glycation end-products, promotes proliferation of human arterial smooth muscle cells 1676
Rippin J.D., Patel A., Belyaev N.D., Gill G.V., Barnett A.H., Bain S.C.: Nitric oxide synthase gene polymorphisms and diabetic nephropathy (*Short Communication*) 426
Ritz-Laser B., Gauthier B.R., Estreicher A., Mamin A., Brun T., Ris F., Salmon P., Halban P.A., Trono D., Philippe J.: Ectopic expression of the beta-cell specific transcription factor Pdx1 inhibits glucagon gene transcription 810
Robertson M.D., Currie J.M., Morgan L.M., Jewell D.P., Frayn K.N.: Prior short-term consumption of resistant starch enhances postprandial insulin sensitivity in healthy subjects 659
Rodríguez-Mañas L., Angulo J., Vallejo S., Peiró C., Sánchez-Ferrer A., Cercas E., López-Dóriga P., Sánchez-Ferrer C.F.: Early and intermediate Amadori glycosylation adducts, oxidative stress, and endothelial dysfunction in the streptozotocin-induced diabetic rats vasculature 556
Rosenbauer J., Herzig P., Giani G.: Atopic eczema in early childhood could be protective against Type 1 diabetes 784
Rudich A., Konrad D., Török D., Ben-Romano R., Huang C., Niu W., Garg R.R., Wijesekara N., Germinario R.J., Bilan P.J., Klip A.: Indinavir uncovers different contributions of GLUT4 and GLUT1 towards glucose uptake in muscle and fat cells and tissues 649
Ryan C.M., Geckle M.O., Orchard T.J.: Cognitive efficiency declines over time in adults with Type 1 diabetes: effects of micro- and macrovascular complications 940
Sadeharju K., Knip M., Hiltunen M., Åkerblom H.K., Hyöty H.: The HLA-DR phenotype modulates the humoral immune response to enterovirus antigens 1100
Saito F., Kawaguchi M., Izumida J., Asakura T., Maehara K., Maruyama Y.: Alteration in haemodynamics and pathological changes in the cardiovascular system during the development of Type 2 diabetes mellitus in OLETF rats 1161
Saraheimo M., Teppo A.-M., Forsblom C., Fagerudd J., Groop P.-H., on behalf of the FinnDiane Study Group: Diabetic nephropathy is associated with low-grade inflammation in Type 1 diabetic patients 1402
Schaeffer G., Levak-Frank S., Spitaler M.M., Fleischhacker E., Esenabhalu V.E., Wagner A.H., Hecker M., Graier W.F.: Intercellular signalling within vascular cells under high D-glucose involves free radical-triggered tyrosine kinase activation 773
Schannwell C.M., Schneppenheim M., Perings S.M., Zimmermann T., Plehn G., Strauer B.E.: Alterations of left ventricular function in women with insulin-dependent diabetes mellitus during pregnancy 267
Schousboe K., Visscher P.M., Henriksen J.E., Hopper J.L., Sørensen T.I.A., Kyvik K.O.: Twin study of genetic and environmental influences on glucose tolerance and indices of insulin sensitivity and secretion 1276
Schulze M.B., Manson J.E., Willett W.C., Hu F.B.: Processed meat intake and incidence of Type 2 diabetes in younger and middle-aged women 1465
Soedamah-Muthu S.S., Chang Y.-F., Otvos J., Evans R.W., Orchard T.J.: Lipoprotein subclass measurements by nuclear magnetic resonance spectroscopy improve the prediction of coronary artery disease in Type 1 Diabetes. A prospective report from the Pittsburgh Epidemiology of Diabetes Complications Study 674
Sparre T., Reusens B., Cherif H., Larsen M.R., Roepstorff P., Fey S.J., Mose Larsen P., Remacle C., Nerup J.: Intrauterine programming of fetal islet gene expression in rats – effects of maternal protein restriction during gestation revealed by proteome analysis 1497
Staiger H., Kausch C., Guirguis A., Weisser M., Maerker E., Stumvoll M., Lammers R., Machicao F., Häring H.-U.: Induction of adiponectin gene expression in human myotubes by an adiponectin-containing HEK293 cell culture supernatant (*Rapid Communication*) 956
Strachan M.W.J., Ewing F.M.E., Frier B.M., McCrimmon R.J., Deary I.J.: Effects of acute hypoglycaemia on audi-

tory information processing in adults with Type I diabetes 97
The DECODE-DECODA Study Group, The European Diabetes Epidemiology Group, The International Diabetes Epidemiology Group: Age, body mass index and Type 2 diabetes – associations modified by ethnicity 1063
The European Nicotinamide Diabetes Intervention Trial (ENDIT) Group, Gale E.A.M.: Intervening before the onset of Type 1 diabetes: baseline data from the European Nicotinamide Diabetes Intervention Trial (ENDIT) 339
Toivonen A.M., Kulmala P., Savola K., Åkerblom H.K., Knip M., the Childhood Diabetes in Finland Study Group: Soluble adhesion molecules in pre-clinical Type 1 diabetes: a prospective study (*Short Communication*) 492
Tschoepe D., Menart B., Ferber P., Altmann C., Haude M., Haastert B., Roesen P.: Genetic variation of the platelet-surface integrin GPIIb-IIIa (*PIA1/A2*-SNP) shows a high association with Type 2 diabetes mellitus 984
Tsimaratos M., Roger F., Chabardès D., Mordasini D., Hasler U., Doucet A., Martin P.-Y., Féraille E.: C-Peptide stimulates Na^+,K^+-ATPase activity via PKC alpha in rat medullary thick ascending limb (Rapid Communication) 124
Veiga-da-Cunha M., Delplanque J., Gillain A., Bonthron D.T., Boutin P., Van Schaftingen E., Froguel P.: Mutations in the glucokinase regulatory protein gene in 2p23 in obese French caucasians 704
Vidal J., Verchere C.B., Andrikopoulos S., Wang F., Hull R.L., Cnop M., Olin K.L., LeBoeuf R.C., O'Brien K.D., Chait A., Kahn S.E.: The effect of apolipoprotein E deficiency on islet amyloid deposition in human islet amyloid polypeptide transgenic mice 71
Walter M., Albert E., Conrad M., Keller E., Hummel M., Ferber K., Barratt B.J., Todd J.A., Ziegler A.-G., Bonifacio E.: IDDM2/insulin VNTR modifies risk conferred by IDDM1/HLA for development of Type 1 diabetes and associated autoimmunity 712
Walter U., Toepfer T., Dittmar K.E.J., Kretschmer K., Lauber J., Weiss S., Servos G., Lechner O., Scherbaum W.A., Bornstein S.R., Boehmer H. von, Buer J.: Pancreatic NOD beta cells express MHC class II protein and the frequency of $I\text{-}A^{g7}$ mRNA-expressing beta cells strongly increases during progression to autoimmune diabetes 1106
Weigert C., Friess U., Brodbeck K., Häring H.U., Schleicher E.D.: Glutamine:fructose-6-phosphate aminotransferase enzyme activity is necessary for the induction of TGF-β1 and fibronectin expression in mesangial cells (*Short Communication*) 852
Wenyi Z., Suzuki S., Hirai M., Hinokio Y., Tanizawa Y., Matsutani A., Satoh J., Oka Y.: Role of urotensin II gene in genetic susceptibility to Type 2 diabetes mellitus in Japanese subjects 972
Williams A.J.K., Norcross A.J., Dix R.J., Gillespie K.M., Gale E.A.M., Bingley P.J.: The prevalence of insulin autoantibodies at the onset of Type 1 diabetes is higher in males than females during adolescence (*Short Communication*) 1354
Williams R.M., Amin R., Shojaee-Moradie F., Umpleby A.M., Acerini C.L., Dunger D.B.: The effects of a specific growth hormone antagonist on overnight insulin requirements and insulin sensitivity in young adults with Type 1 diabetes mellitus 1203
Wolf G., Reinking R., Zahner G., Stahl R.A.K., Shankland S.J.: Erk 1,2 phosphorylates $p27^{Kip1}$: Functional evidence for a role in high glucose-induced hypertrophy of mesangial cells 1090
Yanagisawa K., Hingstrup Larsen L., Andersen G., Drivsholm T., Cederberg A., Westergren R., Borch-Johnsen K., Pedersen O., Enerbäck S., Hansen T.: The *FOXC2*-512C>T variant is associated with hypertriglyceridaemia and increased serum C-peptide in Danish Caucasian glucose-tolerant subjects (*Short Communication*) 1576
Yang W.-S., Hsiung C.A., Ho L.-T., Chen Y.-T., He C.-T., Curb J.D., Grove J., Quertermous T., Chen Y.-D.I., Kuo S.-S., Chuang L.-M., the Sapphire Study Group: Genetic epistasis of adiponectin and *PPARγ2* genotypes in modulation of insulin sensitivity: a family-based association study 977
Zecchin H.G., Bezerra R.M.N., Carvalheira J.B.C., Carvalho-Filho M.A., Metze K., Franchini K.G., Saad M.J.A.: Insulin signalling pathways in aorta and muscle from two animal models of insulin resistance – the obese middle-aged and the spontaneously hypertensive rats 479
Zethelius B., Byberg L., Hales C.N., Lithell H., Berne C.: Proinsulin and acute insulin response independently predict Type 2 diabetes mellitus in men – report from 27 years of follow-up study 20
Zhao J., Yang J., Gregersen H.: Biomechanical and morphometric intestinal remodelling during experimental diabetes in rats 1688
Zhu Q., Yamagata K., Miura A., Shihara N., Horikawa Y., Takeda J., Miyagawa J., Matsuzawa Y.: T130I mutation in *HNF-4α* gene is a loss-of-function mutation in hepatocytes and is associated with late-onset Type 2 diabetes mellitus in Japanese subjects 567
Zitzmann M., Gromoll J., Eckardstein A. von, Nieschlag E.: The CAG repeat polymorphism in the androgen receptor gene modulates body fat mass and serum concentrations of leptin and insulin in men 31

For debates

Hales C.N., Ozanne S.E.: *For Debate*: Fetal and early postnatal growth restriction lead to diabetes, the metabolic syndrome and renal failure 1013
Shafrir E., Raz I.: Diabetes: mellitus or lipidus? 433

Letters

Badenhoop K., Pani M.A., Van Autreve J.E., Gorus F.K., Van der Auwera B.J.: Author's reply 591
Beyan H., Valorani M.G., Pozzilli P.: To: Gale EAM (2002) A missing link in the hygiene hypothesis? Diabetologia 45:588–592 301
Bhatia E., Kordonouri O., Balasubramanium K., Rajeswari J., Landt O., Simon P., Lerch M.M., Witt H.: Absence of association between *SPINK1* trypsin inhibitor mutations and Type 1 or 2 diabetes mellitus in India and Germany 1710
Björntorp P.: – to: Shafrir E, Raz I (2003) *For Debate*. Diabetes: mellitus or lipidus? Diabetologia 46:433–440 1586
Blatter-Garin M.C., Kalix B., De Pree S., James R.W.: Aspirin use is associated with higher serum concentrations of the anti-oxidant enzyme, paraoxonase-1 593
Ciechanowski K., Kędzierska K., Herdzik E., Bober J., Domański L., Borowiak K., Różański J., Myślak M.: Long-term hyperglycaemia decreases vascular fraction of extracellular superoxide dismutase 1026
Elsner M., Tiedge M., Lenzen S.: Mechanism underlying resistance of human pancreatic beta cells against toxicity of streptozotocin and alloxan 1713
Frederiksen L., Brødbaek K., Fenger M., Jørgensen T., Borch-Johnsen K., Madsbad S., Urhammer S.A.: No interactions between polymorphisms in the β_3-adrenergic receptor gene and the PPAR-γ gene on the risk of the insulin resistance syndrome in the Danish MONICA cohort 729
Funk G.-C., Bauer E., Oschatz E., Zauner C., Schneeweiss B.: Compensatory hypochloraemic alkalosis in diabetic ketoacidosis 871
Gale E.: Author's reply 302
Hermann R., Veijola R., Sipilä I., Knip M., Åkerblom H.K., Simell O., Ilonen J.: – to: Pani MA, Van Autreve J, Van Der Auwera BJ, Gorus FK, Badenhoop K (2002) Non-transmitted maternal HLA DQ2 or DQ8 alleles and risk of Type I diabetes in offspring: the importance of foetal or post partum exposure to diabetogenic molecules. Diabetologia 45:1340–1343 588
Hockaday T.D.R., Yajnik C.S.: To: Hales CN, Barker DJP (1992) Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype

hypothesis. Diabetologia 35:595–601 303
Klöting I., Poetsch M., Müller-Myhsok B., Rjasanowski I., Kerner W., Klöting N.: Is there relatedness between maternal lines of Type 1 diabetic patients? 441
Krach K., Pani M.A., Seidl C., Van Autreve J., Van der Auwera B.J., Gorus F.K., Tönjes R.R., Badenhoop K.: *DQ-LTR13* modifies Type 1 diabetes (IDDM) susceptibility on high risk *DQ* haplotypes: reply to the comments of Pascual et al. 870
Lambert A.P., Gillespie K.M., Bingley P.J., Gale E.A.M.: – to: Pani MA, Van Autreve J, Van der Auwera BJ, Gorus FK, Badenhoop K (2002) Non-transmitted maternal HLA DQ2 or DQ8 alleles and risk of Type 1 diabetes in offspring: the importance of foetal or post partum exposure to diabetogenic molecules. Diabetologia 45:1340–1343 590
Love-Gregory L., Wasson J., Lin J., Skolnick G., Suarez B., Permutt M.A.: An E23K single nucleotide polymorphism in the islet ATP-sensitive potassium channel gene (Kir6.2) contributes as much to the risk of Type II diabetes in Caucasians as the PPARγ Pro12Ala variant 136
Maclaren N.K., Lan M.S., Schatz D., Malone J., Notkins A.L., Krischer J.: Multiple autoantibodies as predictors of Type 1 diabetes in a general population 873
Mauvais-Jarvis F., Boudou P., Sobngwi E., Riveline J.P., Kevorkian J.P., Villette J.M., Porcher R., Vexiau P, Gautier J.F.: The polymorphism Gly574Ser in the transcription factor HNF-1α is not a marker of adult-onset ketosis-prone atypical diabetes in Afro-Caribbean patients 728
Meigs J.B., Panhuysen C.I.M., Cupples L.A.: Author's reply 1588
Miura J., Pollin T.I., Hu Y., Harashima S.-I., Notkins A.L., Shuldiner A.R.: Autoantibodies in Type 1 and Type 2 diabetes in the Old Order Amish of Lancaster County, Pennsylvania 1024
Ng D.P.K., Warram J.H., Krolewski A.S.: To: Rippin JD, Patel A, Belyaev ND, Gill GV, Barnett AH, Bain SC (2003) Nitric oxide synthase gene polymorphisms and diabetic nephropathy. Diabetologia 46:426–428 1706
Novials A., Kistauri A., Chico A., Gomis R.: – To: Poa NR, Cooper GJS, Edgar PF: Amylin gene promoter mutations predispose to Type 2 diabetes in New Zealand Maori. Diabetologia 46:574–578 1708
Østergård T., Hansen T.K., Nyholm B., Gravholt C.H., Djurhuus C.B., Hosoda H., Kangawa K., Schmitz O.: Circulating ghrelin concentrations are reduced in healthy offspring of Type 2 diabetic subjects, and are increased in women independent of a family history of Type 2 diabetes 134
Pascual M., Koeleman B.P., Eerligh P., Roep B.O., Martin J.: Distribution of HERV-LTR elements in *HLA-DQB1* alleles: comments on the article by Bieda et al. 869
Populaire C., Mori Y., Dina C., Vasseur F., Vaxillaire M., Kadowaki T., Froguel P.: Does the – 11377 promoter variant of *AMP1* gene contribute to the genetic risk for Type 2 diabetes mellitus in Japanese families? 443
Pouwer F., Adriaanse M.C.: To: M. Koopmanschap: Coping with Type 2 diabetes: the patient's perspective. Diabetologia 45:S18–S22 302
Relimpio F.: Comment 1707
Rotondi M., Romagnani P., Brozzetti A., Santeusanio F., Serio M., Falorni A: – to: Nicoletti F, Conget L, Di Mauro M et al. (2002) Serum concentrations of the interferon-α-inducible chemokine IP-10/CXCL 10 are augmented in both newly-diagnosed Type I diabetes mellitus patients and subjects at risk of developing the disease. Diabetologia 45:1107–1110 1020
Shafrir E., Raz I.: Author's reply 1587
Song S.: To: Saleh MC, Wheeler MD, Chan CB (2002) Uncoupling protein-2: evidence for its function as a metabolic regulator. Diabetologia 45:174–187 132
Svensson J., Carstensen B., Mortensen H.B., Borch-Johnsen K.: Gender-associated differences in Type 1 diabetes risk factors? 442
't Hart L.M., Dekker J.M., Heine R.J., Maassen J.A.: Lack of association between gene variants in the *ALMS1* gene and Type 2 diabetes mellitus 1023
Tilburg J.H.O. van, Wijmenga C., Haeften T.W. van: Comment 1588
Tripathy D., Aljada A., Dandona P.: Free Fatty acids (FFA) and endothelial dysfunction; role of increased oxidative stress and inflammation. To: Steinberg et al. (2002) Vascular function, insulin resistance and fatty acids 300
Visser J., Groen H., Klatter F., Vis L., Strubbe J., Rozing J.: The diabetes prone BB rat model of IDDM shows duration of breastfeeding to influence Type 1 diabetes development later in life 1711
Weedon M.N., Gloyn A.L., Davey Smith G., Ben-Shlomo Y., Frayling T.M., Hattersley A.T.: Quantitative traits associated with the Type 2 diabetes susceptibility allele in Kir6.2 1021
Zychma M.J., Gumprecht J., Rutkowski P., Grzeszczak W.: — to: Rippin JD, Patel A, Belyaev ND, Gill GV, Barnett AH, Bain SC (2003) Nitric oxide synthase gene polymorphisms and diabetic nephropathy. Diabetologia 46:426–428 1707

Book reviews

138, 446, 1028

EASD News section

R 1, R 11, R 13, R 15, R 17, R 27, R 41, R 47, R 51, R 55, R 67, R 77

Errata

140, 1296, 1446, 1589

Supplement 1

Ceriello A.: The possible role of postprandial hyperglycaemia in the pathogenesis of diabetic complications M9
Del Prato S.: Loss of early insulin secretion leads to postprandial hyperglycaemia M2
Dunning B.E., Deacon R., Gutierrez C., Paladini S., Valentin M.A., Foley J.E.: Pharmacologic restoration of the early insulin response in pre-diabetic monkeys controls mealtime glucose excursions without peripheral hyperinsulinaemia M22
Hu S., Boettcher B.R., Dunning B.E.: The mechanisms underlying the unique pharmacodynamics of nateglinide M37
Qiao Q., Tuomilehto J., Borch-Johnsen K.: Post-challenge hyperglycaemia is associated with premature death and macrovascular complications M17
Standl E., Füchtenbusch M.: The role of oral antidiabetic agents: why and when to use an early-phase insulin secretion agent in Type 2 diabetes mellitus M30
Van Gaal L.F., De Leeuw I.H.: Rationale and options for combination therapy in the treatment of Type 2 diabetes M44

Supplement 2

Abstracts of the 18th Congress of the International Diabetes Federation Paris, France, 24–29 August 2003 A1

List of referees

1715

Author index

1720

Contents of Volume 46

I